


AD-A280 464



DOCUMENTATION PAGE

Form Approved
OMB No 0704-0188

It is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and reviewing the collection of information, sending comments regarding this burden estimate or any other aspect of this burden estimate, to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 1993		3. REPORT TYPE AND DATES COVERED Reprint	
4. TITLE AND SUBTITLE (see title on reprint)				5. FUNDING NUMBERS PE: NWED QAXM WU: 00129	
6. AUTHOR(S) Rokita et al.					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Armed Forces Radiobiology Research Institute 8901 Wisconsin Ave. Bethesda, MD 20889-5603				8. PERFORMING ORGANIZATION REPORT NUMBER SR94-6	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Uniformed Services University of the Health Sciences 4301 Jones Bridge Road Bethesda, MD 20814-4799				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.					
13. ABSTRACT (Maximum 200 words) DTIC ELECTE JUN 20 1994 S B D DTIC QUALITY INSPECTED 3					
14. SUBJECT TERMS				15. NUMBER OF PAGES 5	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED		18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED		19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	
				20. LIMITATION OF ABSTRACT	

INCREASED FIBRINOGEN SYNTHESIS IN MICE DURING THE ACUTE PHASE RESPONSE: CO- OPERATIVE INTERACTION OF INTERLEUKIN 1, INTERLEUKIN 6, AND INTERLEUKIN 1 RECEPTOR ANTAGONIST

Hanna Rokita,* Ruta Neta,† Jean D. Sipe‡

Interleukin 6 (IL-6) stimulates fibrinogen (Fg) gene expression both in vivo and in vitro; while interleukin 1 (IL-1) paradoxically stimulates in vivo, yet inhibits in vitro, Fg synthesis. The naturally occurring interleukin 1 receptor antagonist (IL-1ra) and passive immunization with anti-IL-6 antiserum were used to study the in vivo mechanism of action of IL-1 on Fg gene expression. Changes in plasma Fg and hepatic Fg mRNA concentrations were measured following administration of exogenous IL-1ra together with IL-6 or IL-1 to CD2F1 mice. Our results suggest that in vivo, IL-1 per se inhibits Fg production since when IL-1ra was co-administered with IL-6, greater concentrations of Fg were observed than when IL-6 was administered alone. The data suggest that IL-1 stimulates Fg production through intermediate production of IL-6, since stimulation was abrogated when either IL-1ra or anti-IL-6 antiserum was co-administered with IL-1. An in vivo role for IL-1ra in the stimulation of Fg by IL-1 was supported by the observation that within 1 h of IL-1 administration to mice, IL-1ra mRNA was detectable in liver. It appears that IL-1, an early mediator of inflammation, inhibits constitutive expression of Fg genes and stimulates the IL-1ra and IL-6 genes. The inhibitory effect of IL-1 is reversed by endogenous IL-1ra and by the direct stimulation of Fg gene expression by IL-6.

Changes in hepatic protein synthesis are a major manifestation of the acute phase response and are mainly regulated by the mononuclear phagocyte system through secretion of cytokines such as interleukin 6 (IL-6) and interleukin 1 (IL-1).¹⁻⁶ IL-6 and, to a lesser extent, IL-1 have been implicated as modulators of hepatic serum amyloid A (SAA) and fibrinogen (Fg) mRNA and/or protein synthesis in humans and mice.⁴⁻⁷ Treatment of mice with endotoxin and the two lipopolysaccharide-induced cytokines, IL-1 and TNF, caused an increase in plasma Fg concen-

tration.⁸ IL-1 is an early and pluripotent mediator of the acute phase response, with capacity to induce itself, IL-6 and a structurally related receptor ligand with antagonist activity, termed IL-1 receptor antagonist (IL-1ra). Recently, recombinant generated preparations of IL-1ra, a 17 kDa protein with 26-30% homology to IL-1 beta and 19% to IL-1 alpha have been used to distinguish physiological roles of IL-1 from closely related cytokines such as IL-6.⁹⁻¹⁷ Since several in vitro studies^{5,18-21} indicate that constitutive and IL-6 stimulated Fg gene expression in hepatoma cell lines is inhibited by IL-1, the present study was undertaken to investigate the basis of the in vivo stimulatory effect of IL-1 on Fg gene expression.

From the *Institute of Molecular Biology, The Jagiellonian University, Krakow, Poland; †Department of Experimental Hematology, Armed Forces Radiobiology Research Institute, Bethesda, MD 20889, USA; and ‡Department of Biochemistry, Boston University School of Medicine, Boston, MA 02118, USA

Correspondence to: Dr Jean D. Sipe, Department of Biochemistry, Boston University School of Medicine, K121, 80 East Concord Street, Boston, MA 02118, USA

Received 15 January 1993; revised and accepted for publication 12 March 1993

© 1993 Academic Press Limited
1043-4666/93/050454-05 \$08.00/0

KEY WORDS: Fibrinogen IL-1 IL-6 IL-1ra Serum amyloid A

RESULTS

IL-1ra Inhibits IL-1, but Enhances IL-6, Stimulation of Plasma Fibrinogen

In vivo administration of exogenous cytokines demonstrates biological activity and permits com-



parison of dose response and sensitivity to individual cytokines. Although IL-1 alpha and IL-6 both stimulated a concentration-related elevation in Fg, IL-1 was more effective on a weight basis than IL-6 (Table 1).

TABLE 1. Stimulation of in vivo fibrinogen production by cytokines.

	Treatment			Fibrinogen (% control)
	IL-1 (ng)	IL-6 (μg)	IL-1ra (μg)	
1.	-	-	-	100
2.	10	-	-	147 ± 15
3.	300	-	-	264 ± 5
4.	-	0.25	-	146
5.	-	1.25	-	157 ± 20
6.	300	-	100	115 ± 15
7.	-	-	75	110
8.	-	1.25	10	167 ± 7
9.	-	1.25	75	194 ± 7
10.	-	1.25	100	213 ± 23

Mice were bled 24 h after injections. Values are means and standard errors of two to three experiments, each three or four mice per group. Exceptions are numbers 4 and 7 where data from one experiment were available.

The stimulatory effect of IL-1 on Fg production was blocked by the co-administration of IL-1ra with IL-1. In contrast, the stimulation of Fg by IL-6 was enhanced by co-administration of IL-1ra and IL-6.

Endogenous IL-1ra Production in Response to IL-1 but Not IL-6

Endogenous expression of IL-1ra mRNA in liver was detectable 1 and 1.5 h after administration of IL-1 alone and in combination with IL-6, but not after IL-6 alone (Fig. 1). The polyadenylated RNA enriched




Figure 1. Kinetics of IL-1ra mRNA, relative to beta-actin mRNA in livers of mice.

Results are given 0, 1, 1.5 and 5 h after administration of IL-1 (---▲---; 300 ng), IL-6 (—■—; 1.25 μg) alone or in combination (—●—) to mice. Northern blots were prepared from mouse liver RNA enriched in polyadenylated mRNA, 3–5 μg/lane. Results are representative of four experiments.

fractions contained varying amounts of ribosomal RNA, leading to fluctuation in the relative abundance of beta-actin mRNA. Therefore the results shown in Fig. 1 are normalized for the content of beta actin in each preparation: the quantities of beta-actin and GAPDH mRNA were comparable. IL-1ra mRNA was not detectable at 30 min, appeared to reach maximal levels at 2–3 h and was expressed at a diminished level by 5.5 h after IL-1 administration.

IL-1 Stimulates Fg and SAA Production by Induction of IL-6

IL-6 has been shown to mediate other in vivo responses to IL-1;^{22–23} the role of IL-6 in IL-1 stimulated Fg gene expression was indicated by the reduction in stimulation of Fg after administration of 20 ng IL-1 to mice passively immunized with anti-IL-6 antiserum 16 h earlier (Table 2) as compared with Fg

TABLE 2. Passive immunization with anti-IL-6 antibodies blocks stimulation of fibrinogen by IL-1.

Pretreatment	Fibrinogen		
	Protein (% control)	mRNA (Fg Actin mRNA)	SAA (μg ml)
None	176 ± 15	1.0	248 ± 25
Anti-IL-6	118 ± 17	0.6	93 ± 4
Rat Ig Control	310 ± 15	nd	196 ± 2
GL13, Ab Control	182 ± 9	1.2	220 ± 16

Each of three mice per group were injected with 600 μg of each Ig 16–20 h prior to stimulation with 20 ng of IL-1. Values are mean and standard errors of individual plasma samples obtained 24 h after administration of IL-1; nd, not done. Results are representative of two to four experiments.

concentration in mice that received no pretreatment or received control immunoglobulins. The ratio of hepatic Fg mRNA to hepatic actin mRNA 1 h after administration of IL-1 to mice pretreated with anti-IL-6 antibodies was 50% of that in mice treated with control immunoglobulins or in mice with no pretreatment (Fig. 2). The in vivo production of IL-6 following IL-1 was further demonstrated by reduction in plasma SAA (Table 2) which is known to be produced in response to the synergistic action of IL-1 and IL-6.²⁴ We used a hybridization probe for gamma-Fg in these studies; expression of alpha, beta and gamma chains has been shown to proceed essentially coordinately²⁵ and in our study hybridization with either gamma-Fg or alpha-Fg specific cDNA probes revealed similar kinetics (Figs 1, 2).




Figure 2. Kinetics of Fg and IL-1ra (mRNA, relative to beta-actin mRNA in livers of mice.

Results are given 0, 1, 1.5 and 5 h after administration of IL-1 (20 ng), IL-6 (1.25 μ g) alone and in combination to mice that received anti-IL-6 antibodies (600 μ g), control immunoglobulins (600 μ g) or no treatment 16 h earlier. — Δ —, IL-1; — \square —, IL-6; — \blacksquare —, IL-1 + IL-6; — \times —, IL-1 + anti IL-6. Northern blots were prepared from mouse liver RNA enriched in polyadenylated mRNA, 3–5 μ g/lane. Insert shows the Northern hybridization signal for the 1 h time point. Results are representative of four experiments.

DISCUSSION

This study has addressed the conflicting observations that IL-1 suppresses Fg gene expression in hepatoma cell lines, yet enhances Fg gene expression in mice.^{1,2} The basis for increased concentrations of plasma Fg during the acute phase response, has, for the first time, been investigated by combining an analysis of plasma protein changes with changes in mRNA expression in liver. Using IL-1ra together with IL-6 we can confirm in the *in vivo* situation the previously reported *in vitro* inhibition of Fg gene expression by IL-1. Our results also provide strong evidence that, *in vivo*, IL-1 counteracts its direct inhibitory effect on Fg gene expression by stimulation of IL-1ra and IL-6 gene expression. In turn, IL-6 directly stimulates Fg gene expression and IL-1ra blocks inhibition of Fg gene expression by IL-1 leading to increased Fg production in the presence of IL-6.

The observation of IL-1ra mRNA expression in mice following IL-1 but not IL-6 administration (Fig. 2) suggests that IL-1ra exerts both temporal and bidirectional quantitative regulation of acute phase changes in liver protein synthesis by counteracting

both the synergistic interaction of IL-1 with IL-6 required for SAA induction and its antagonistic effect on Fg induction by IL-6.^{5,18–22,24} IL-1 has been shown to influence the effects of IL-6 on plasma proteins other than Fg. For example, IL-1 acts synergistically with IL-6 to enhance SAA gene expression by Hep 3B and Hep G2 cells.²⁴ IL-1 has been shown to decrease the stimulatory effect on IL-6 on alpha-2-macroglobulin as well as Fg in Hep G2 cells.⁵ These reports together with our findings in this paper indicate that IL-1 and IL-6 interactions may be one of the factors that contribute to the unique kinetic changes that individual acute phase reactants undergo during the acute phase response.

The mechanism by which IL-1 reduces constitutive Fg production and IL-6 induced Fg gene up-regulation is not understood. The Fg genes are expressed constitutively in liver and are further stimulated by dexamethasone and IL-6 and indirectly by IL-1 and TNF.^{1,3,22,26} The studies of Huber and co-workers demonstrated the interplay between IL-6 and glucocorticoid responsive elements in the promoter region of Fg genes.²⁶ Fluctuation in receptor ratios and/or occupancy may play a role in the inhibitory effect of IL-1 on Fg gene expression. The recent studies of Nesbitt and Fuller²⁷ indicate that IL-1 and, to a lesser extent, IL-6, may elicit a transient reduction in the number of IL-6 receptors in rat hepatocytes, whereas IL-6 and IL-1 increased IL-6 receptor mRNA in human primary hepatocytes.²⁸ Other explanations may lie in the selective interaction of transcription factors. Thus, while our study has resolved the paradox of the conflicting *in vivo* and *in vitro* effects of IL-1 on Fg gene expression, the question as to how IL-1 down-regulates Fg remains.

MATERIALS AND METHODS

Experimental Animals

Female CD2F1-mice, 6–8 weeks of age, free from overt disease, were employed. All animal handling procedures were performed as described²⁹ in compliance with guidelines from the National Research Council and the Armed Forces Radiobiology Research Institute. At various time points, citrated plasma was collected and livers were removed and snap-frozen in liquid nitrogen and stored at -85°C for subsequent mRNA analysis.

Reagents

Recombinant human IL-1 alpha was the gift of Dr Peter Lomedico of Hoffman-LaRoche, Nutley, NJ. The preparation, lot IL-1 2/88 was stored in aliquots at -20°C until use. Recombinant human interleukin 6, lot PPG9001 was provided by Dr E. Liehl, Sandoz Pharme AG, Basel.

Switzerland. The IL-1ra was provided by Dr Robert Thompson, Syngen, Boulder, CO. Rat monoclonal antibody to recombinant mouse IL-6 (MP5 20F3) was prepared using partially purified Cos-7 mouse IL-6 as immunogen (22). Rat monoclonal antibody to beta-galactosidase was used as an isotype control.²³

Protein Assays

Fibrinogen in citrated plasma was measured as the rate of conversion of fibrinogen to fibrin in the presence of excess thrombin using the Sigma Diagnostic Kit (Sigma Chemical Co., St Louis, MO) as aibrator to express the data as milligrams of fibrinogen per 100ml plasma. Measurements of fibrin clot formation were performed on a fibrometer (Becton-Dickinson Co., Fairleigh, N.J.). Mouse SAA was measured by ELISA (Hemagen Diagnostics, Waltham, MA).

RNA Extraction and Northern Blot Hybridization Analysis

Polyadenylated RNA was extracted from individual and pooled livers using Mini RiboSep mRNA isolation kits (Collaborative Research, Bedford, MA) according to the manufacturer's instructions. RNA was denatured by glyoxylation and size-fractionated by electrophoresis through 1.5% agarose, and transferred to Genescreen Plus nylon membranes (Dupont NEN, Boston, MA). The quantity of IL-1ra,¹⁶ fibrinogen,³⁰ actin³¹ and GAPDH³² mRNA in each lane was determined by hybridization with the corresponding cDNA probes labelled with alpha-32P-dCTP (Dupont NEN, Boston, MA) using a random priming oligonucleotide labelling kit (Bethesda Research Laboratories, Gaithersburg, MD). Unincorporated cCTP was removed from the reaction mixture through the use of Nensorb columns (Dupont NEN). Following overnight hybridization at 42°C, filters were washed as previously described³³ and exposed to X-ray film at -70°C for 2-24h. The relative abundance of SAA mRNA was measured relative to beta actin by scanning densitometry of autoradiographs following Northern blot hybridization.

Acknowledgements

We thank Drs Ivan Otterness and Aleksander Koj for helpful discussions and Faith Selzer and Isabel Carreras for assistance with experiments. This work was supported in part by U.S. Public Health Service Grant AG9006 and by grant 2173 from Pfizer, Inc. This work was supported by the Armed Forces Radiobiology Research Institute, Defense Nuclear Agency, under work unit 00129. Views presented in this paper are those of the authors; no endorsement by the Defense Nuclear Agency or the Department of Defense has been given or should be inferred. Research was conducted according to the principles

enunciated in the Guide for the Care and Use of Laboratory Animals, prepared by the Institute of Laboratory Animal Resources, National Research Council.

REFERENCES

1. Gordon AH, Koj A (1985) The Acute Phase Response to Injury and Infection. Elsevier, Amsterdam, pp 1-339.
2. Sipe JD (1989) The molecular biology of interleukin-1 and the acute phase response. *Adv Internal Med* 34:1-20.
3. Baumann H, Gauldie J (1990) Regulation of hepatic acute phase plasma protein genes by hepatocyte stimulating factors and other mediators of inflammation. *Mol Biol Med* 7:147-159.
4. Magielska-Zero D, Bereta J, Czuba-Pelech B, Pajdak W, Gauldie J, Koj A (1988) Inhibitory effect of recombinant human interferon gamma on synthesis of some acute phase proteins in human hepatoma HepG2 stimulated by leukocyte cytokines, TNF alpha and INF-beta2/BSF2/IL-6. *Biochem Int* 17:17-23.
5. Darlington GJ (1989) Response of liver genes to acute phase mediators. *Ann NY Acad Sci* 557:310-315.
6. Neta R, Vogel SN, Sipe JD, Wong GG, Nordan RP (1988) Comparison of in vivo effects of human recombinant IL-1 and human recombinant IL-6 in mice. *Lymphokine Res* 7:403-412.
7. Rokita H, Szuba K (1991) Regulation of acute phase reaction by transforming growth factor beta in cultured murine hepatocytes. *Acta Biochim Polon* 38:241-249.
8. Bertini R, Bianchi M, Erroi A, Villa P, Ghezzi P (1989) Dexamethasone modulation of in vivo effects of endotoxin, tumor necrosis factor, and interleukin-1 on liver cytochrome P-450, plasma fibrinogen, and serum iron. *J Leuk Biol* 46:254-262.
9. Arend WP (1991) Interleukin-1 receptor antagonist: A new member of the interleukin 1 family. *J Clin Invest* 88:1445-1451.
10. Eisenberg SP, Evans RJ, Arend WP, Verdeber E, Brewer MT, Hannum CH, Thompson RC (1990) Primary structure and functional expression from complementary DNA to a human interleukin-1 receptor antagonist. *Nature* 343:341-346.
11. Carter DB, Deibel MR, Dunn CJ, Tomich C-SC, Laborde AL, Slightom AL, Berger AE, Bienkowski MJ, Sun FF, McEwan RN, Harris PKW, Yem AW, Waszak GA, Chosay JG, Sieu LC, Hardee MM, Zurcher-Neely HA, Reardon IM, Heinrichson RL, Truesdell SE, Shelly JA, Eessalu TE, Taylor BM, Tracey DE (1990) Purification, cloning, expression and biological characterization of an interleukin-1 receptor antagonist protein. *Nature* 344:633-638.
12. Vogel SN, Henricson BE, Neta R (1991) Roles of interleukin-1 and tumor necrosis factor in lipopolysaccharide-induced hypoglycemia. *Infect Immun* 59:2494-2498.
13. Henricson BE, Neta R, Vogel SN (1991) Interleukin-1 receptor antagonist blocks lipopolysaccharide-induced colony-stimulating factor production and early endotoxin tolerance. *Infect Immun* 59:1188-1191.
14. Neta R, Abrams J, Perlstein R, Rokita H, Sipe JD, Vogel SN, Whitnall M (1992) Interaction of cytokines, IL-1, TNF and IL-6 in in vivo host responses. In Revel M (ed.), IL-6, Pathophysiology and Clinical Potential, Raven Press, New York, pp 195-202.
15. Numerof R, Sipe JD, Zhang Z, Dinarello CA, Mier JW (1990) Use of the IL-1 receptor antagonist (IL-1ra) in IL-2 treated mice. *Lymphokine Res* 9:612.
16. Zahedi K, Seldin MF, Rits ME, Ezekowitz AB, Whitehead AS (1991) Mouse IL-1 receptor antagonist protein. Molecular characterization, gene mapping, and expression of mRNA in vitro and in vivo. *J Immunol* 146:4228-4233.

17. McIntyre KW, Stepan GJ, Kolinsky KD, Benjamin WR, Plocinski JM, Kaffka KL, Campen CA, Chizzonite RA, Kilian PL (1991) Inhibition of interleukin-1 (IL-1) binding and bioactivity in vitro and modulation of acute inflammation in vivo by IL-1 receptor antagonist and anti-IL-1 receptor monoclonal antibody. *J Exp Med* 173:931-939.
18. Mackiewicz A, Speroff T, Ganapathi MK, Kushner I (1991) Effects of cytokine combinations on acute phase protein production in two human hepatoma cell lines. *J Immunol* 146:3032-3037.
19. Malawista SE, Van Damme J, Sipe JD, Duff GW, Weiss MC (1989) Independent effects of interleukin 6 (IL-6) and interleukin 1 (IL-1) on accumulation of specific mRNA for fibrinogen and for serum amyloid A in human hepatoma (HepG2) cells. *Ann NY Acad Sci* 557:518-520.
20. Darlington GJ, Wilson DR, Lachman LB (1986) Monocyte-conditioned medium, interleukin-1 and tumor necrosis factor stimulate the acute phase response in human hepatoma cell lines. *J Cell Biol* 103:787-793.
21. Heinrich P, Castell JV, Andus T (1990) Interleukin-6 and the acute phase response. *Biochem J* 265:621-636.
22. Starnes HE, Pearce MK, Tewari A, Yim JH, Zou JC, Abrams JS (1990) Anti-IL-6 monoclonal antibodies protect against lethal *Escherichia coli* infection and lethal tumor necrosis factor challenge in mice. *J Immunol* 145:4185-4191.
23. Neta R, Perlstein R, Vogel SN, Ledney GD, Abrams J (1992) Role of interleukin 6 (IL-6) in protection from lethal irradiation and in endocrine responses to IL-1 and tumor necrosis factor. *J Exp Med* 175:689-694.
24. Sipe JD, Rokita H, Bartle LM, Loose LD, Neta R (1991) The IL-1 receptor antagonist simultaneously inhibits SAA and stimulates fibrinogen synthesis in vivo and in vitro: a proposed mechanism of action. *Cytokine* 3:497.
25. Fuller GM, Otto JM, Woloski BM, McCarty CL, Adams MA (1985) The effects of hepatocyte stimulating factor on fibrinogen biosynthesis in hepatocyte monolayers. *J Cell Biol* 101:1481-1486.
26. Huber P, Laurent M, Dalmon J (1990) Human beta-fibrinogen gene expression: Upstream sequences involved in its tissue specific expression and its dexamethasone and interleukin 6 stimulation. *J Biol Chem* 265:5695-5701.
27. Nesbitt JL, Fuller GM (1992) Differential regulation of interleukin-6 receptor and gp130 gene expression in rat hepatocytes. *Mol Biol Cell* 3:103-112.
28. Bauer J, Bauer TM, Kalb T, Jaga T, Lengyel G, Hirano T, Kishimoto T, Acs G, Mayer L, Gerok W (1989) Regulation of interleukin-6 receptor expression in human monocytes and monocyte-derived macrophages. *J Exp Med* 170:1537-1549.
29. Perlstein RS, Mougey EH, Jackson WE, Neta R (1991) Interleukin-1 and interleukin-6 act synergistically to stimulate the release of adrenocorticotrophic hormone in vivo. *Lymphokine Cytokine Res* 10:141-146.
30. Bolvard MG, Lord SI (1988) High-level expression of a functional human fibrinogen gamma chain in *Escherichia coli*. *Gene* 66:183-192.
31. Bond JF, Farmer SE (1983) Regulation of tublin and actin mRNA production in rat brain: Expression of a new beta tubulin mRNA with development. *Mol Cell Biol* 3:1333-1342.
32. Tso JY, Sun X-H, Kao T-h, Reece KS, Wu R (1985) Isolation and characterization of rat and human glyceraldehyde-3-phosphate dehydrogenase cDNAs: genomic complexity and molecular evolution of the gene. *Nucleic Acids Res* 13:2485-2502.
33. Rokita H, Shirahama T, Cohen AS, Meek RL, Benditt EP, Sipe JD (1987) Differential expression of the amyloid SAA3 gene in liver and peritoneal macrophages of mice undergoing dissimilar inflammatory episodes. *J Immunol* 139:3849-3853.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DIIC TAB	<input type="checkbox"/>
Chem Abstracts	<input type="checkbox"/>
Indexing/Classification	
By	
Distribution	
Availability Order	
Avail and/or	
Dist Special	
A-1	20